

LITHIUM IN THE UPPER-CENTAURUS LUPUS AND LOWER-CENTAURUS CRUX SUBGROUPS OF SCORPIUS-CENTAURUS

ERIC J. BUBAR¹

Department of Biology and Physical Sciences, Marymount University, 2807 N. Glebe Road, Arlington, VA 22207

MARC SCHAEUBLE, JEREMY R. KING

Department of Physics and Astronomy, Clemson University, Clemson, SC 29630-0978

ERIC E. MAMAJEK

Department of Physics and Astronomy, University of Rochester, P.O. Box 270171, Rochester, NY 14627-0171

JOHN R. STAUFFER

Spitzer Science Center, Caltech, Pasadena, CA 91125

Draft version October 31, 2011

ABSTRACT

We utilize spectroscopically derived model atmosphere parameters and the Li I $\lambda 6104$ subordinate line and the $\lambda 6708$ doublet to derive lithium abundances for 12 members of the Upper-Centaurus Lupus (UCL) and Lower-Centaurus Crux (LCC) subgroups of the Scorpius Centaurus OB Association. The results indicate any intrinsic Li scatter in our 0.9-1.4 M_{\odot} stars is limited to ~ 0.15 dex, consistent with the lack of dispersion in $\geq 1.0 M_{\odot}$ stars in the 100 Myr Pleiades and 30-50 Myr IC 2391 and 2602 clusters. Both ab initio uncertainty estimates and the derived abundances themselves indicate that the $\lambda 6104$ line yields abundances with equivalent or less scatter than is found from the $\lambda 6708$ doublet as a result of lower uncertainties for the subordinate feature, a result of low sensitivity to broadening in the subordinate feature. Because NLTE corrections are less susceptible to changes in surface gravity and/or metallicity for the 6104 Å line, the subordinate Li feature is preferred for deriving lithium abundances in young Li-rich stellar association stars with $T_{\text{eff}} \geq 5200$ K. At these temperatures, we find no difference between the Li abundances derived from the two Li I lines. For cooler stars, having temperatures at which main-sequence dwarfs show abundance patterns indicating overexcitation and overionization, the $\lambda 6104$ -based Li abundances are ~ 0.4 dex lower than those derived from the $\lambda 6708$ doublet. The trends of the abundances from each feature with T_{eff} suggest that this difference is due to (near)UV photoionization, which in NLTE preferentially ionizes Li atoms in the subordinate $2p$ state relative to $2s$ resonance line state due to opacity effects. Consequently, this overionization of Li in the $2p$ state, apparently not adequately accounted for in NLTE corrections, weakens the $\lambda 6104$ feature in cooler stars. Accordingly, the $\lambda 6708$ -based abundances may give more reliable estimates of the mean Li abundance in cool young stars. Our mean Li abundance, $\log N(\text{Li}) = 3.50 \pm 0.07$ is ~ 0.2 dex larger than the meteoritic value. While stellar models suggest that Li depletion of at least 0.4 dex, and possibly much larger, should have occurred in our lowest mass star(s), our Li abundances show no decline with decreasing mass indicative of such depletion.

Subject headings: stars: abundances - stars: late-type

1. INTRODUCTION

The identification of the lower mass members of Scorpius-Centaurus, the nearest OB Association to our Sun, has provided a rich source of nearby (within 150 pc) sun-like pre-MS stars for understanding early stellar evolution. Sco-Cen comprises three primary subgroups: Upper-Sco (US), Upper-Centaurus Lupus (UCL) and Lower-Centaurus Crux (LCC) with ages of ~ 10 Myr (Pecaut et al. 2011), ~ 15 Myr and ~ 16 Myr (Preibisch & Mamajek 2008), respectively. One of the key pieces of the puzzle in understanding young stars and their internal structure lies within understanding their lithium abundances. Since lithium is depleted during

the pre-main sequence (by 0.30-1.0 dex for a 1 M_{\odot} , solar composition star (Piau & Turck-Chieze 2002)), it provides a strong constraint on internal stellar structure and has provided a valuable diagnostic of youth, yet several puzzling questions remain regarding lithium depletion.

Historically, lithium has been a valuable target for much abundance work, particularly regarding open clusters. Stars with $M > 1 M_{\odot}$ in the 100 Myr Pleiades cluster and in the 30-50 Myr IC 2391 and 2602 clusters evince no Li star-to-star Li dispersion above the uncertainties (Figures 3 and 7 of King et al. 2000; Figure 4 of Randich et al. 2001). One of the most vexing problems lies in the star-to-star dispersion in lithium abundances for $M \leq 0.8 M_{\odot}$ in the Pleiades cluster (Soderblom 1993; King et al. 2010), a result at odds with standard stellar models and the lack of abundance scatter in lower mass IC 2391 and

¹ Department of Physics and Astronomy, University of Rochester, P.O. Box 270171, Rochester, NY 14627-0171

2602 stars. In addition Li dispersion of the size seen in the Pleiades is not apparent in lower mass stars in clusters of older ages (e.g. Hyades: Soderblom et al. 1995).

Russell (1996) examined this Pleiades dispersion by exploring lithium abundances derived from two lithium features, the strong doublet at 6708 Å ($2p$ - $2s$ transition) and the weaker subordinate feature at 6104 Å ($3d$ - $2p$ triplet transition) (Carlsson et al. 1994, Grotrian diagram in Figure 3 of). He found that abundances from the subordinate line were greater than those from the resonance doublet and that less scatter was observed in lithium abundances derived from equivalent widths of the 6104 Å line (± 0.1 dex) when compared with those derived from the 6708 Å line (± 0.2 dex). Russell (1996) suggested that a larger-scale survey of lithium in the Pleiades using the 6104 Å feature might explain the Pleiades Li dispersion as an artifact of spurious $\lambda 6708$ -based abundances.

Ford et al. (2002) reexamined the work of Russell (1996) using spectra of 11 late-G and early-K Pleiads and suggested that improper accounting of an Fe II blend in the 6104 Å region was likely responsible for both the decreased spread and enhanced abundances they reported. By utilizing spectral synthesis of the 6104 Å which accounted for this Fe II blending feature, the spread in abundances derived by Ford et al. (2002) was equal for both the $\lambda 6104$ and $\lambda 6708$ lines, indicating a genuine abundance spread for the Pleiades. In addition, while some abundances from the subordinate line agreed with those from the 6708 doublet, there were 4 stars which showed clearly larger abundances from $\lambda 6104$, which Ford et al. (2002) suggested might be the result of spot coverage on those stars.

Most recently, King et al. (2010) revisited lithium in the Pleiades and compared lithium abundances derived from both $\lambda 6104$ and $\lambda 6708$. They also found that the scatter in 6104 Å abundances persisted and was comparable to that observed in the 6708 Å results. However, contrary to Ford et al. (2002), they found little difference between abundances from the two lines; the source of these discrepant conclusions is unclear. Salient questions that remain are which line gives the most reliable Li abundances, and whether the Pleiades Li scatter is a relic of differential PMS depletion already in place by the ~ 15 Myr age of our stars. While our sample's mass range limits its utility for exploring the latter, it is well-suited for former.

Recent work suggests that ZAMS-to-Hyades age dwarfs with $T_{\text{eff}} \leq 5200$ K show abundance anomalies which mimic the effects of overionization (Schuler et al. 2003). The reality of overionization in such stars can be explored using the two Li I features alone by exploiting the NLTE elements of Li I line formation at such temperatures. Due to a conspiracy of the photoionizing energies and the UV line opacity, the 6104 Å lower energy level ($2p$) can be depopulated relative to the 6708 Å ground state ($2s$) (Carlsson et al. 1994). If the extent of this effect is not accounted for in NLTE calculations, the $\lambda 6104$ feature, weakened relative to the resonance line, would yield lower abundances. Might the Li-rich stars in Sco-Cen yield 6104 Å lines strong enough to measure abundances of sufficient accuracy to search for such an

offset?

Motivated by these questions, we have derived lithium for a sample of UCL and LCC members of Sco-Cen. Being young (~ 15 Myr), the lithium abundances in our stars are necessarily large, which decreases the uncertainty in spectral synthesis of the weaker 6104 Å lines which may affect the aforementioned studies of the Pleiades. For our mass range, the expected modest depletion factors allow us to determine abundances through spectral synthesis of both the $\lambda 6104$ feature and the $\lambda 6708$ doublet. We therefore derive lithium for a sample of 12 UCL and LCC members from these two lithium features. While our mass range limits the viability of answering questions regarding depletion mechanisms, we are able to robustly constrain the lithium abundances of our stars using the traditionally weaker $\lambda 6104$ subordinate line as well as the $\lambda 6708$ doublet. We compare the results from the two lines, analyze the differences in terms of potential NLTE effects, make recommendations for which features to use in determining lithium abundances in other young clusters and associations, and determine a mean abundance of lithium for the UCL/LCC subgroups of Sco-Cen.

2. DATA, OBSERVATIONS AND ANALYSIS

2.1. Spectroscopic Observations and Reductions

The sample explored here is a subset of the sample of Mamajek et al. (2002), who identified members of the UCL and LCC subgroups of the Scorpius-Centaurus OB Association using x-ray, proper-motions and color-magnitude selection. Optical spectroscopy of the sample was obtained on June 14-17, 2002 with the CTIO 4 meter Blanco telescope and the echelle spectrograph with 31.6 l/mm grating and a 2048X2048 CCD detector. The 0.8" slit width yielded a resolution of $R \sim 40,000$ with a typical S/N of 60-80 per summed pixel in exposure times of ≈ 100 -300 s. The spectra have incomplete wavelength coverage extending from approximately 5800 Å to 7800 Å, across 45 orders. The spectra have been reduced using standard routines in the echelle package of IRAF². These include bias correction, flat-fielding, scattered light correction, order extraction, and wavelength calibration. We have selected stars which meet three criteria for this study: 1) spectra are higher S/N (~ 60 -80) 2) stars are apparently slower rotators ($v \sin i \leq 20$ km s⁻¹) and 3) spectra showed no evidence of multiplicity in their cross-correlation peaks when correlated with a template solar spectrum taken with the same instrument. Further details of the velocity and multiplicity results can be found in Bubar et al. (2011).

3. ABUNDANCE ANALYSIS AND UNCERTAINTIES

Abundances from the Li I resonance feature at 6708 Å and the 1.8 eV subordinate features at 6104 Å were derived from spectral synthesis using an updated version of the LTE analysis package MOOG (Sneden 1973). Stellar parameters were derived using excitation and ionization balance following the approach described in Bubar & King (2010) and are listed in Ta-

² IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

ble 1. Model atmospheres corresponding to these parameters were interpolated from the Kurucz ATLAS9 grids³. The $\lambda 6104$ linelist was that described and used by King et al. (2010). The $\lambda 6708$ region linelist was that used by King et al. (1997) updated using the CN features from Mandell, Ge & Murray (2004) and additional atomic data from Kurucz⁴ and VALD (Piskunov et al. 1995; Kupka et al. 1999).

For MML 28, comparison of the synthesis with the observed data in the 6708 Å region suggested a differential radial velocity shift between Li and other metal lines that could not be rectified with variations in smoothing or input abundances in a fashion that yielded acceptable fits to the 6104 Å region. The inferred differential shift of the 6708 Å feature was blueward by 0.7 km/s (−0.016 Å). Cayrel et al. (2007) found that 3D NLTE modeling of the Li I feature yielded a redshifted line with respect to 1D LTE results, inconsistent with the shift we observed in MML 28. Given the doublet structure of the $\lambda 6708$ feature, it is possible that the magnetic intensification mechanism described by Leone (2007) could strengthen the blue component of the resonance line relative to the red component. However, our results indicate that MML 28 does not evince any line strength enhancement, relative to other stars, that should accompany such a wavelength shift relative to other stars in the sample. There is, however, observational evidence (Allende Prieto & García López 1998; Reddy et al. 2002; Mandell, Ge & Murray 2004) that convective motions can produce Li I resonance lines with blueshifted centroids. In the 6708 Å synthesis for MML 28, we have simply shifted the synthetic Li lines by −0.016 Å, roughly 2–3 times the shift of the stronger Li hyperfine components applied by Mandell, Ge & Murray (2004) in their linelist calibrated with the Sun.

Input abundances for the initial syntheses were solar values scaled according to the [Fe/H] for each star derived by Bubar et al. (2011). Small adjustments (a couple hundredths of a dex) within the uncertainties of Bubar et al. (2011) were allowed if they consistently improved the fitting of blending features neighboring both the 6104 and 6708 Å Li features. Smoothing was accomplished utilizing a Gaussian with a fixed (in each wavelength region) FWHM corresponding to the instrumental resolution ($R \sim 40,000$) and then employing rotational broadening which consistently yielded the best fit to the line profiles in both the 6104 and 6708 Å region. As expected, the $v \sin i$ values we determined in this fashion are (except for MML 30 and MML 73) slightly smaller than the values measured by Bubar et al. (2011) that parameterized the total broadening (instrumental, thermal, rotational, etc).

Representative observed spectra and syntheses (including our best fit values) are shown in Figures 1 and 2 for the 6104 and 6708 Å regions respectively. Comparison of our spectra with syntheses having no Li clearly and confidently indicates the presence of the blended, moderate strength $\lambda 6104$ Li I feature in all but one of our stars’ spectrum (MML 73). For this star, our sense is that the $\lambda 6104$ is present and measurable, but 2 uncoop-

erative pixels in the midst of the blended Li feature do not make this conclusion definitive (see Figure 3). Our $\lambda 6104$ abundance is included in all of the statistical analyses (unless stated otherwise), but the star is denoted as an inverted triangle in the figures. The best fit LTE Li abundances are provided in columns 2 and 4 of Table 2.

NLTE corrections were applied to both the $\lambda 6104$ and $\lambda 6708$ values according to the prescription of Carlsson et al. (1994). Interpolations within their grid of NLTE corrections provided the necessary correction to our LTE lithium abundances, based on T_{eff} , $\log g$, [Fe/H], and $\log N(\text{Li})$. Three stars had absolute physical parameters which placed them outside the range of values populated by the grid. Within the adopted uncertainties of the parameters, however, the stars all fall within an acceptable extrapolation range. For MML 72, which had a supersolar metallicity of 0.06, we adopted a correction which assumed solar metallicity, justified based on the apparent insensitivity of the grid to changes in [Fe/H] (e.g. changes in metallicity of −0.71 dex and −0.14 dex are required to alter the 6104 and 6708 Å NLTE corrections by 0.01 dex at the T_{eff} , $\log g$ and Li abundances characterizing the star). Both MML 43 and MML 44 had surface gravities which were outside of the range covered by the Carlsson et al. (1994) grid. For these stars we assumed a surface gravity of 4.5, the maximum value in the grids. The sensitivity of the NLTE corrections to changes in the surface gravity is small: a −0.20 dex difference in gravity for both MML 43 and MML 44 results in a $\leq +0.01$ dex difference in the NLTE corrections for the 6708 Å line, and $\log g$ changes of ≥ 1.0 dex are needed to observe a 0.01 dex change in NLTE corrections for the 6104 Å line). This suggests that surface gravity changes have a negligible effect on the NLTE corrections, given the uncertainties. Final NLTE Li abundances are provided in columns 3 and 5 of Table 2.

Uncertainties in the Li abundances are dominated by two general components: uncertainties in the fitting and the T_{eff} values (uncertainties in $\log g$ and ξ contribute negligibly). The fitting uncertainties are dominated by uncertainties in the continuum location and $v \sin i$ values. Plausible necessarily subjective estimates of allowed variations in these quantities were made and the Li abundances rederived to estimate the fitting uncertainty. These fitting uncertainties were added in quadrature with the abundance uncertainties due to the uncertainty in T_{eff} from Bubar et al. (2011) to estimate the total uncertainty in our Li abundance from each of the 6104 and 6708 Å features. These are listed in columns 3 and 5 of Table 2. We note that the uncertainties in the $\lambda 6104$ - and $\lambda 6708$ -based Li abundances due to T_{eff} uncertainty are correlated. When we later examine the difference between the Li abundances derived from the two wavelength regions, we account for the correlation of this component of the uncertainties in calculating the uncertainty of the difference.

4. RESULTS AND DISCUSSION

4.1. Lithium Dispersion

The 6104 Å- and 6708 Å-based NLTE Li abundances are plotted versus T_{eff} in Figure 4. In traditional open cluster work, the T_{eff} coordinate serves as a monotonic

Fig1
Fig2
Fig3
Tab2

³ <http://kurucz.harvard.edu/grids.html>

⁴ <http://kurucz.harvard.edu>

mass coordinate, but this does not hold in very young star formation regions/associations such as Sco-Cen. Accordingly, we plot the NLTE Li abundances versus mass in Figure 5. We utilized bolometric luminosities from Mamajek et al. (2002) and spectroscopic effective temperatures to interpolate within the pre-main sequence grids of D’Antona & Mazzitelli (1997) in order to derive masses for our stars. It should be noted that our track-based masses may be underpredicted. According to Hillenbrand & White (2004), masses from evolutionary tracks underpredict PMS stellar masses by as much as 10-30 % when compared with dynamical masses. Such an offset could presumably account for some of the lack of observed depletion, but quantifying this is difficult with currently available models. Consequently we proceed with the analysis making use of our derived masses.

The ordinary Pearson correlation coefficients indicate no significant variation in either Li abundance with stellar mass. A striking result is the pleasingly small scatter in the $\lambda 6104$ Li I abundances compared to the $\lambda 6708$ -based values. The smaller scatter in the former is consistent with the smaller uncertainties we find for the former. Historically, the $\lambda 6104$ line has not frequently been utilized in abundance studies due to its weakness and increased blending compared to the $\lambda 6708$ resonance feature. Our results, however, suggest that the subordinate feature may produce higher quality abundances than the resonance feature in young Li-rich stars. In addition, the NLTE corrections are significantly less sensitive to changes in both metallicity and surface gravity for the 6104 line. Independent changes in $\log g$ of $\sim \pm 0.20$ dex and $\sim \pm 0.15$ dex in $[\text{Fe}/\text{H}]$ yield changes of $\sim \pm 0.01$ dex in the NLTE corrections for the 6708 line. A change of order 1.0 dex and 0.71 in $\log g$ and $[\text{Fe}/\text{H}]$, respectively, is needed to see a 0.01 dex change in the NLTE correction for the 6104 line. Based on the lower scatter from the 6104 Å line and the lower sensitivity to parameter changes ($\log g$ and $[\text{Fe}/\text{H}]$) for NLTE corrections to the subordinate line, we conclude, similarly to Carlsson et al. (1994) (see Figure 10 of that work), that the 6104 Å feature is more well-suited for obtaining lithium abundances in young, lithium rich solar-type stars.

The rms scatter in the $\lambda 6104$ - $\lambda 6708$ -based Li abundances about their respective means in Figure 5 (0.15 and 0.25 dex, respectively) compares favorably with the typical uncertainties (0.08 and 0.18 dex, respectively). The $\lambda 6708$ -based scatter is dominated by the lower abundance of MML 73; though nothing else appears remarkable or notably different about this star, the $\lambda 6708$ -based rms scatter of 0.17 dex found when excluding it is equivalent to that expected on the basis of the uncertainties alone. We conclude from these comparisons that any real Li dispersion—measured by the square root of the difference between the variance of the Li abundance data and the square of typical Li abundance uncertainty—in our sample of modestly rotating stars is limited to ≤ 0.15 dex.

The $\lambda 6708$ - and $\lambda 6104$ -based Li abundance of cool slowly-rotating Pleiads with $T_{\text{eff}} \leq 5500$ K, which characterizes most of the stars in our Sco-Cen sample, show a real dispersion of ~ 0.6 dex (King et al. 2010). Comparable spreads in $\lambda 6708$ -based abundances are seen for $T_{\text{eff}} \leq 5200$ K in the 50-90 Myr α Per cluster

(Balachandran, Lambert & Stauffer 1996; Stauffer et al. 1999; Balachandran et al. 2011). However, mass is the relevant variable in considering PMS Li depletion, and the masses of the Pleiads that evince significant dispersion are some 0.3-0.6 M_{\odot} lower in mass than the stars in our sample. Pleiads having masses comparable to those objects in our sample show no significant Li dispersion (e.g., Figure 4 of King, Krishnamurthi & Pinsonneault 2000). Our data therefore do not address the interesting question of whether the Pleiades and α Per Li dispersion is the relic of possible differential Li depletion mechanisms having acted at the ~ 15 Myr age of Sco-Cen.

4.2. $\lambda 6104$ versus $\lambda 6708$

Figure 6 shows the difference between the NLTE $\lambda 6708$ - and $\lambda 6104$ -based Li abundances versus T_{eff} . The mean difference (6708 minus 6104) is $+0.20 \pm 0.07$ (mean uncertainty) dex. The 7 stars with $T_{\text{eff}} \geq 5200$ K show no significant difference in the Li abundances derived from the two Li features, a result consistent with the abundances derived in similarly warmer Pleiads (King et al. 2010). Figure 6 indicates that the stars with statistically significant 0.3-0.5 dex abundance differences are the coolest objects in our sample.

Fig6

The behavior of this difference may qualitatively fit the picture of magnetic intensification suggested by Leone (2007), where magnetic fields of 10^{-1} to 1 T can enhance the strength of strong Li I resonance features relative to weak ones. Whether such an effect would: a) hold for the $\lambda 6104$ feature, (b) hold for stars of our T_{eff} , gravity, and Li abundance, and (c) would “switch on” at $T_{\text{eff}} \leq 5200$ K as we observe, all require additional calculations beyond those carried out by Leone (2007).

Recent works point to the existence of abundance patterns consistent with overexcitation/ionization in cool ($T_{\text{eff}} \leq 5200$ K) open cluster dwarfs (Schuler et al. 2003, 2004, 2006). However, such a mechanism (whatever its cause[s]) acting in the framework of LTE can not explain the difference and the sense of the difference we observe in our coolest stars. Figure 2 provides an important clue to the source of the behavior in demonstrating that the $\lambda 6708$ - $\lambda 6104$ difference is due to declining $\lambda 6104$ abundances (and not increasing $\lambda 6708$ abundances) with declining T_{eff} . We believe that this is a signature of relative NLTE overphotoionization at low T_{eff} acting to selectively depopulate the $2p$ level (the lower level of the $\lambda 6104$ feature relative to the $2s$ level (the lower level of the $\lambda 6708$ feature), thus weakening the $\lambda 6104$ line relative to the $\lambda 6708$ line. Indeed, Carlsson et al. (1994) note the importance of ultraviolet photoionization as a NLTE mechanism important for Li I line formation and PMS stars exhibit greater UV emission (Findeisen & Hillenbrand 2010). A significant finding of their work is that, at low T_{eff} and high Li abundance, (near)ultraviolet metal line opacity inhibits photoionization from the Li $2s$ level, allowing photoionization from the $2p$ level to dominate.

Our results and those of King et al. (2010) indicate that the relative $\lambda 6104$ versus $\lambda 6708$ Li I NLTE corrections from Carlsson et al. (1994) are adequate for young Li rich stars with $T_{\text{eff}} \geq 5200$ K, but the $\lambda 6104$ corrections are too small, perhaps by ≥ 0.4 dex, at cooler T_{eff} . This suggests that while the internal uncertainties and sensitivity of NLTE corrections to stellar parameters is

TABLE 1
PHYSICAL PARAMETERS

Name	Teff K	logg	[Fe/H]	MtVel km s ⁻¹	log($\frac{L}{L_{\odot}}$) dex	Mass M _⊙
MML 7	5763 ± 132	4.24 ± 0.34	-0.01 ± 0.11	2.54 ± 0.14	0.39 ± 0.06	1.3
MML 13	4984 ± 125	3.99 ± 0.54	-0.27 ± 0.10	2.53 ± 0.28	0.26 ± 0.10	1.5
MML 28	4975 ± 82	4.08 ± 0.40	-0.21 ± 0.05	2.08 ± 0.16	-0.37 ± 0.14	0.9
MML 30	5180 ± 68	4.02 ± 0.30	-0.13 ± 0.05	2.18 ± 0.12	-0.05 ± 0.09	1.1
MML 36	5198 ± 115	4.39 ± 0.29	-0.09 ± 0.09	1.96 ± 0.22	0.00 ± 0.05	1.2
MML 40	5440 ± 113	4.36 ± 0.20	-0.22 ± 0.09	3.16 ± 0.22	0.07 ± 0.08	1.2
MML 43	5580 ± 98	4.60 ± 0.28	-0.07 ± 0.08	2.59 ± 0.25	0.06 ± 0.08	1.1
MML 44	5526 ± 145	4.71 ± 0.36	-0.05 ± 0.12	2.29 ± 0.29	0.54 ± 0.10	1.5
MML 55	5228 ± 80	4.28 ± 0.24	-0.12 ± 0.07	2.50 ± 0.16	0.16 ± 0.11	1.3
MML 70	5040 ± 100	4.09 ± 0.20	-0.18 ± 0.10	2.24 ± 0.15	0.01 ± 0.10	1.2
MML 72	4950 ± 125	4.19 ± 0.44	0.06 ± 0.13	1.31 ± 0.42	0.21 ± 0.07	1.2
MML 73	5212 ± 97	4.00 ± 0.32	-0.20 ± 0.09	2.42 ± 0.19	0.21 ± 0.01	1.4

smaller for the $\lambda 6104$ feature relative to the $\lambda 6708$ feature, the absolute abundances from the $\lambda 6708$ may be more reliable modulo other possible (perhaps activity-related) effects on the resonance line (King & Schuler 2004).

4.3. Sco-Cen Lithium Abundance

Since the $\lambda 6104$ Li I abundances are likely not free of systematic errors in the NLTE corrections for the cooler stars as discussed above, our mean Li abundance estimate of $\log N(\text{Li}) = 3.50 \pm 0.08$ (mean uncertainty) is made from the $\lambda 6708$ -based NLTE results. This abundance is $\sim 2\sigma$ larger than the meteoritic value. While no mass-dependent slope to the Li data indicative of depletion is seen in Figure 5, even the most conservative PMS Li depletion models predict depletion factors of ≥ 0.4 dex for $1 M_{\odot}$ stars at an age of 15 Myr (Piau & Turck-Chieze 2002) (see their Figure 4). Our results suggest the initial

Li abundance of our cooler objects might be a factor of 2-3 higher than observed today. On the other hand, the existence of PMS Li depletion inhibiting mechanisms remains possible (Ventura et al. 1998). Assessing whether the degree of Li depletion suffered by our Sco-Cen sample and whether our data are in agreement with PMS Li depletion models will require extending the Li-mass relation via observation and analyses of cooler, lower mass Sco-Cen members that are expected to be more prodigious depleters.

EJB and EEM acknowledge support from NSF grant AST 10-08908 to EEM and the University of Rochester School of Arts and Sciences. MS and JRK gratefully acknowledge support for this work from NSF grant AST 09-08342 to JRK. We would also like to acknowledge the referee whose comments helped to clarify the work.

REFERENCES

- Allende Prieto, C., & Garcia Lopez, R. J. 1998, *A&AS*, 129, 41
Balachandran, S., Lambert, D. L., & Stauffer, J. R. 1996, *ApJ*, 470, 1243
Balachandran, S. C., Mallik, S. V., & Lambert, D. L. 2011, *MNRAS*, 410, 2526
Bubar, E. J., & King, J. R. 2010, *AJ*, 140, 293
Bubar, E. J., Peca, M. J., Mamajek, E. E., & Bourque, M. 2011, *ApJ*, in preparation
Carlsson, M., Rutten, R. J., Bruls, J. H. M. J., & Shchukina, N. G. 1994, *A&A*, 288, 860
Cayrel, R., et al. 2007, *A&A*, 473, L37
D’Antona, F., & Mazzitelli, I. 1997, *Mem. Soc. Astron. Italiana*, 68, 807
D’Orazi, V., & Randich, S. 2009, *A&A*, 501, 553
Findeisen, K., & Hillenbrand, L. 2010, *AJ*, 139, 1338
Ford, A., Jeffries, R. D., & Smalley, B. 2002, *A&A*, 391, 253
Gustafsson, B., Bell, R. A., Eriksson, K., & Nordlund, A. 1975, *A&A*, 42, 407
Hillenbrand, L. A., & White, R. J. 2004, *ApJ*, 604, 741
King, J. R., Schuler, S. C., Hobbs, L. M., & Pinsonneault, M. H. 2010, *ApJ*, 710, 1610
King, J. R., Deliyannis, C. P., Hiltgen, D. D., Stephens, A., Cunha, K., & Boesgaard, A. M. 1997, *AJ*, 113, 1871
King, J. R., Krishnamurthi, A., & Pinsonneault, M. H. 2000, *AJ*, 119, 859
King, J. R., & Schuler, S. C. 2004, *AJ*, 128, 2898
Kupka, F., Piskunov, N., Ryabchikova, T. A., Stempels, H. C., & Weiss, W. W. 1999, *A&AS*, 138, 119
Leone, F. 2007, *ApJ*, 667, L175
Mamajek, E. E., Meyer, M. R., & Liebert, J. 2002, *AJ*, 124, 1670
Mandell, A. M., Ge, J., & Murray, N. 2004, *AJ*, 127, 1147
Peca, M., Mamajek, E. E., & Mamajek, E. 2011, in preparation
Piau, L., & Turck-Chieze, S. 2002, *ApJ*, 566, 419
Piskunov, N. E., Kupka, F., Ryabchikova, T. A., Weiss, W. W., & Jeffery, C. S. 1995, *A&AS*, 112, 525
Preibisch, T., & Mamajek, E. 2008, *Handbook of Star Forming Regions*, Volume II, 235
Randich, S., Pallavicini, R., Meola, G., Stauffer, J. R., & Balachandran, S. C. 2001, *A&A*, 372, 862
Reddy, B., Lambert, D., Laws, C., Gonzalez, G., & Covey, K. 2002, *MNRAS*, 335, 1005
Russell, S. C. 1996, *ApJ*, 463, 593
Schuler, S. C., King, J. R., Fischer, D. A., Soderblom, D. R., & Jones, B. F. 2003, *AJ*, 125, 2085
Schuler, S. C., King, J. R., Hobbs, L. M., & Pinsonneault, M. H. 2004, *ApJ*, 602, L117
Schuler, S. C., King, J. R., Terndrup, D. M., Pinsonneault, M. H., Murray, N., & Hobbs, L. M. 2006, *ApJ*, 636, 432
Sestito, P., Palla, F., & Randich, S. 2008, *A&A*, 487, 965
Snedden, C. 1973, *ApJ*, 184, 839
Soderblom, D. R., Jones, B. F., Balachandran, S., Stauffer, J. R., Duncan, D. K., Fedele, S. B., & Hudon, J. D. 1993, *AJ*, 106, 1059
Soderblom, D. R., Jones, B. F., Stauffer, J. R., & Chaboyer, B. 1995, *AJ*, 110, 729
Stauffer, J. R., et al. 1999, *ApJ*, 527, 219
Ventura, P., Zepieri, A., Mazzitelli, I., & D’Antona, F. 1998, *A&A*, 331, 1011




FIG. 1.— Sample lithium syntheses of the 6104 Å subordinate line of lithium for MML 30 (top) and MML 43 (bottom). The observed spectrum is plotted as solid squares and three syntheses are shown for each star.




FIG. 2.— Sample lithium syntheses of the 6708 Å doublet of Li for MML 30 (top) and MML 43 (bottom). The observed spectrum is plotted as solid squares and three lithium syntheses are shown for each star.




FIG. 3.— Sample lithium syntheses of the $\lambda 6104$ line of lithium (top) and the $\lambda 6708$ doublet (bottom) for MML 73. This measurable abundance of lithium from the 6104 Å feature is less definitive in this star than in the rest of our sample. The 6708 Å results, however, show the clear presence of Li in the star.




FIG. 4.— The NLTE corrected abundance of lithium is plotted versus temperature for the stars based on the $\lambda 6104$ subordinate feature (top) and the $\lambda 6708$ resonance doublet (bottom). The least squares fit performed in each case is shown as the dotted line, and the ordinary Pearson correlation coefficient between Li abundance in T_{eff} is labeled in the plots.




FIG. 5.— The NLTE abundances, $\log N(\text{Li})_{6104}$ (top) and $\log N(\text{Li})_{6708}$ (bottom), are plotted versus mass. NLTE corrected abundances for a sample of Pleiades stars (from Soderblom 1993) are plotted for $\log N(\text{Li})_{6708}$ (bottom-open squares) as a comparison sample which exhibits depletion. Note that our lowest mass star clearly shows no evidence of depletion, which is contrary to stellar lithium depletion models given the age and assumed mass.




FIG. 6.— The difference in NLTE abundances, $\log N(\text{Li})_{6708} - \log N(\text{Li})_{6104}$ is plotted versus temperature. The difference that is seen at $T_{\text{eff}} \leq 5200$ K is due to the effects of overphotoionization on λ 6104 which manifests itself by yielding lower abundances when compared with the λ 6708-based abundances.

TABLE 2
LITHIUM ABUNDANCE

Name	$\log N(\text{Li})_{6104}$ LTE	$\log N(\text{Li})_{6104}$ NLTE	$\log N(\text{Li})_{6708}$ LTE	$\log N(\text{Li})_{6708}$ NLTE
MML 7	3.47	3.53 ± 0.09	4.22	3.65 ± 0.16
MML 13	3.28	3.43 ± 0.10	4.15	3.84 ± 0.21
MML 28	3.08	3.24 ± 0.08	3.93	3.56 ± 0.13
MML 30	3.18	3.31 ± 0.05	3.98	3.57 ± 0.22
MML 36	3.11	3.23 ± 0.08	4.14	3.79 ± 0.18
MML 40	3.37	3.46 ± 0.08	4.04	3.57 ± 0.15
MML 43	3.37	3.45 ± 0.08	3.88	3.43 ± 0.24
MML 44	3.24	3.32 ± 0.10	3.84	3.40 ± 0.20
MML 55	3.27	3.39 ± 0.07	3.83	3.45 ± 0.18
MML 70	3.20	3.35 ± 0.09	4.07	3.73 ± 0.16
MML 72	2.81	2.97 ± 0.10	3.59	3.29 ± 0.22
MML 73	≤ 2.98	3.11 ± 0.07	3.03	2.86 ± 0.14